SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Rabbit Anti-CD21 antibody

Catalog Number: SLM-60246R

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Monoclonal
Clone No.:	CD12B9
Target:	CD21
Immunogen:	KLH conjugated synthetic peptide derived from human CD21
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Applications:	WB=1:500-2000 IHC-P=1:100-500 IHC-F=1:100-500 ICC/IF=1:100-500 IF=1:100-500 Flow-Cyt=1μg/Test （Paraffin sections need antigen repair） not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	Human
Predicted MW.:	111kDa
Subcellular Location:	The nucleus cytoplasmic
Background:	This gene encodes a membrane protein, which functions as a receptor for Epstein-Barr virus (EBV) binding on B and T lymphocytes. Genetic variations in this gene are associated with susceptibility to systemic lupus erythematosus type 9 (SLEB9). Alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Sep 2009] **Database links:** **Entrez Gene:** 1380 Human **SwissProt:** P20023 Human

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com

[VALIDATION IMAGES]




Figure 1. 25 ug total protein per lane of various lysates (see on figure) probed with CD21 monoclonal antibody, unconjugated (SLM-60246R) at 1:2000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min




Figure 2. Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD21 Monoclonal Antibody, Unconjugated(SLM-60246R) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining




Figure 3. Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD21 Monoclonal Antibody, Unconjugated(SLM-60246R) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining




Figure 4. Paraformaldehyde-fixed, paraffin embedded Human appendix; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD21 Monoclonal Antibody, Unconjugated(SLM-60246R) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining




Figure 5. Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CD21 Monoclonal Antibody, Unconjugated (SLM-60246R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (Green, SL0295G-BF488), DAPI (blue, C02-04002) was used to stain the cell nuclei




Figure 6. Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CD21 Monoclonal Antibody, Unconjugated (SLM-60246R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (Green, SL0295G-BF488), DAPI (blue, C02-04002) was used to stain the cell nuclei

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com




Figure 7. The Raji (H) cells were incubated in 5%BSA to block non-specific protein-protein interactions (30 min at r.t.), followed by secondary antibody incubation for 40 min at room temperature. Primary Antibody (green): Rabbit Anti-CD21 antibody (SLM-60246R): 1 µg/10^6 cells; Isotype Control (orange): Rabbit IgG (SL0295P). Blank control (black): PBS. Acquisition of 20,000 events was performed




Figure 8. 4% Paraformaldehyde-fixed Raji (H) cell; Antibody incubation with (CD21) monoclonal Antibody, unconjugated (SLM-60246R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, SL60295G-BF488) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.