

Rabbit Anti-LMO4 antibody

SLM-60672R

Product Name	LMO4
Chinese Name	乳腺 Tumour 自身抗原 LMO4Recombinant rabbit monoclonal anti LMO 4; LMO-4; Breast tumor autoantigen; LIM domain only 4; LIM domain only protein 4; LIM domain transcription factor LMO 4; LIM domain transcription factor LMO4; LMO4_HUMAN.
Alias	
Immunogen Species	Rabbit
Clonality	Monoclonal
Clone NO.	R9B1
React Species	Human(predicted:Mouse) WB=1:500-2000
Applications	not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Theoretical molecular weight	18kDa
Form	Liquid
Concentration	1mg/ml
immunogen	KLH conjugated synthetic peptide derived from human LMO4
Lsotype	IgG
Purification	affinity purified by Protein A
Buffer Solution	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Attention	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
PubMed	PubMed This gene encodes a cysteine-rich protein that contains two LIM domains but lacks a DNA-binding homeodomain. The encoded protein may play a role as a transcriptional regulator or as an oncogene. [provided by RefSeq, Aug 2008]
Product Detail	Subcellular Location: Nucleus.

SWISS:
P61968

Gene ID:
8543

Database links:


[Entrez Gene: 8543](#) Human

[Entrez Gene: 16911](#) Mouse

[SwissProt: P61968](#) Human

[SwissProt: P61969](#) Mouse

Product Picture



Blocking buffer: 5% NFDN/TBST

Primary Ab dilution: 1:1000

Primary Ab incubation condition: 2 hours at
room temperature

Secondary Ab: Goat Anti-Rabbit IgG H&L

(HRP)



Lysate: 1: HeLa, 2: MCF-7, 3: NIH/3T3

Protein loading quantity: 20 μ g

Exposure time: 60 s

Predicted MW: 18 kDa

Observed MW: 18 kDa