

# Active Brain Derived Neurotrophic Factor (BDNF) Instruction Manual

## SBPA007Mu01

**Mus musculus (Mouse)**

**Buffer Formulation**

20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

**Traits**

Freeze-dried powder

**Purity**

> 90%

**Isoelectric Point**

8.5

**Applications**

Cell culture; Activity Assays.

### ACTIVITY TEST




Figure. The binding activity of BDNF with APP.

Brain-derived neurotrophic factor, also known as BDNF, is a member of the neurotrophin family of growth factors, which are related to the canonical Nerve Growth Factor. BDNF acts on certain neurons of the central nervous system and the peripheral nervous system, helping to support the survival of existing neurons, and encourage the growth and differentiation of new neurons and synapses. Besides, Amyloid Precursor Protein (APP) has been identified as an interactor of BDNF, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse BDNF and recombinant mouse APP.

Briefly, BDNF were diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 $\mu$ L were then transferred to APP-coated microtiter wells and incubated for

2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-BDNF pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of of BDNF and APP was shown in Figure 1, and this effect was in a dose dependent manner.

## USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## Image




Figure. Gene Sequencing (Extract)

## Image




Figure. SDS-PAGE




Figure. Western Blot

**[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.