

Active Calpain 3 (CAPN3) Instruction Manual

SBPC285Hu01

Homo sapiens (Human)

Buffer Formulation	PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.
Traits	Freeze-dried powder
Purity	> 95%
Isoelectric Point	5.5
Applications	Cell culture; Activity Assays.

ACTIVITY TEST




Figure. The binding activity of CAPN3 with TTN.

Calpain 3 is a calcium-dependent cysteine protease mainly expressed in skeletal muscle. In humans, calpain 3 is encoded by the CAPN3 gene. This gene encodes a muscle-specific member of the calpain large subunit family that specifically binds to titin. Mutations in this gene are associated with limb-girdle muscular dystrophies type 2A. Alternate promoters and alternative splicing result in multiple transcript variants encoding different isoforms and some variants are ubiquitously expressed. Besides, Titin (TTN) has been identified as an interactor of CAPN3, thus a binding ELISA assay was conducted to detect the interaction of recombinant human CAPN3 and recombinant human TTN. Briefly, CAPN3 were diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to TTN-coated microtiter wells and

incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-CAPN3 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of CAPN3 and TTN was shown in Figure 1, and this effect was in a dose dependent manner.

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE




Figure. Western Blot

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.