

# Active Fibroblast Growth Factor 13 (FGF13) Instruction Manual

**SBPC322Ra01**

**Rattus norvegicus (Rat)**

**Buffer Formulation**

20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

**Traits**

Freeze-dried powder

**Purity**

> 97%

**Isoelectric Point**

9.2

**Applications**

Cell culture; Activity Assays.

**ACTIVITY TEST**



A




B

**Figure 1. Cell proliferation of 3T3 cells after stimulated with FGF13.**

**(A) 3T3 cells cultured in DMEM, stimulated with 100ng/mL FGF13 for 48h;**

**(B) Unstimulated 3T3 cells cultured in DMEM for 48h.**



**Figure 2. Cell proliferation of 3T3 cells after stimulated with FGF13.**

FGF13 (Fibroblast growth factor 13) is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth, and invasion. A proliferation assay was conducted to detect the bioactivity of recombinant rat FGF13 using 3T3 cells. Briefly, 3T3 cells were seeded into triplicate wells of 96-well plates at a density of 2,000 cells/well and allowed to attach overnight, then the medium was replaced with serum-free standard DMEM prior to the addition of various concentrations of FGF13. After incubated for 48h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 $\mu$ L of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37°C. Proliferation of 3T3 cells after incubation with FGF13 for 48h observed by inverted microscope was shown in Figure 1. Cell viability was assessed by CCK-8 (Cell Counting Kit-8 ) assay after incubation with recombinant FGF13 for 48h. The result was shown in Figure 2. It was obvious that FGF13 significantly increased cell viability of 3T3 cells.

## USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### Image



SDS-PAGE Image




Figure. Western Blot; Sample: Recombinant FGF13, Rat.

### **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

