Eukaryotic Dickkopf Related Protein 1 (DKK1) Instruction Manual

SFPA167Hu61

Homo sapiens (Human)

Source	Eukaryotic expression
Host	CHO Cell
Endotoxin Level	<1.0EU per 1μg (determined by the LAL method)
Subcellular Location	Secreted
Predicted Molecular Mass	27.2kDa
Accurate Molecular Mass	34-44kDa(Analysis of differences refer to the manual)
Residues & Tags	Leu33~His266 with N-terminal His Tag
Buffer Formulation	PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 95%
Isoelectric Point	8.4
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
                                    LNSVLNSN AIKNLPPPLG
GAAGHPGSAV SAAPGILYPG GNKYQTIDNY QPYPCAEDEE CGTDEYCASP
TRGGDAGVQI CLACRKRRKR CMRHAMCCPG NYCKNGICVS SDQNHFRGEI
EETITESFGN DHSTLDGYSR RTTLSSKMYH TKGQEGSVCL RSSDCASGLC
CARHFWSKIC KPVLKEGQVC TKHRRKGSHG LEIFQRCYCG EGLSCRIQKD
HHQASNSSRL HTCQRH
```

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.