

Eukaryotic Stem Cell Factor Receptor (SCFR) Instruction Manual

SFPA391Hu61

Homo sapiens (Human)

Source	Eukaryotic expression
Host	293F cell
Endotoxin Level	<1.0EU per 1 μ g (determined by the LAL method)
Subcellular Location	Membrane, Cytoplasm
Predicted Molecular Mass	57.6kDa
Accurate Molecular Mass	65kDa(Analysis of differences refer to the manual)
Residues & Tags	Gln26~Pro524 with N-terminal His Tag
Buffer Formulation	PBS, pH7.4, containing 5% Trehalose.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	6.4
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
QPSVS PGEPSPPSIH PGKSDLIVRV
GDEIRLLCTD PGFVKWTFEI LDETNNENKQN EWITEKAEAT NTGKYTCTNK
HGLSNSIYVF VRDPAKLFLV DRSLYGKEDN DTLVRCPLTD PEVTNYSLKG
CQ GKPLPKDL RFIPDPKAGI MIKSVKRAYH RLCLHCSVDQ EGKSVLSEKF
ILKVRPAFKA VPVVS VKAS YLLREGEEFT VTCTIKDVSS SVYSTWKREN
SQT KLQEKYN SWHHGDFNYE RQATLTISSA RVNDSGVFMC YANNTFGSAN
VTTTLEVVDK GFINIFPMIN TTVFVNDGEN VDLIVEYEAF PKPEHQQWIY
MNRFTDKWE DYPKSENESEN IRYVSELHLT RLKGTEGGTY TFLVSNSDVN
AAIAFNVVVN TKPEILTYDR LVNGMLQVA AGFPEPTIDW YFCPGTEQRC
SASVLPVDVQ TLNSSGPPFG KLVVQSSIDS SAFKHNGTVE CKAYNDVGKT
SAYFNFAFKG NNKEQIHPHT LFTP
```

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.