

Eukaryotic Immunoglobulin G1 (IgG1) Instruction Manual

SFPA402Rb61

Oryctolagus cuniculus (Rabbit)

Source	Eukaryotic expression
Host	293F cell
Endotoxin Level	<1.0EU per 1ug (determined by the LAL method)
Subcellular Location	Secreted
Predicted Molecular Mass	25kDa
Accurate Molecular Mass	33kDa(Analysis of differences refer to the manual)
Residues & Tags	Glu1~Lys180 with N-terminal His Tag and C-terminal Fc Region of Rabbit IgG
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 97%
Isoelectric Point	8.2
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
GQPKAPSVFP LAPCCGDTPS STVTLGCLVK GYLPEPVTVT WNSGTLTNGV  
RTFPSVRQSS GLYLSSSVVS VTSSSQPVTC NVAHPATNTK VDKTVAPSTC  
SKPTCPPPEL LGGPSVFIFP PKPKDTLMIS RTPEVTCVVV DVSQDDPEVQ  
FTWYINNEQV RTARPPLREQ QFNSTIRVVS
```

USAGE

Reconstitute in PBS or others.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.