

Eukaryotic Coagulation Factor XII (F12) Instruction Manual

SFPA622Ra61

Rattus norvegicus (Rat)

Source	Eukaryotic expression
Host	293F cell
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Secreted
Predicted Molecular Mass	27.6kDa
Accurate Molecular Mass	34kDa(Analysis of differences refer to the manual)
Residues & Tags	Val354~Phe595 with N-terminal His Tag
Buffer Formulation	PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	5.2
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
VVGGGLVA LPGSHPYIAA LYWGDSFCAG SLIDPCWVLT AAHCLQKRPA  
PEELTVVLGQ DRHNQSCERC QTLAVHSYRL HEGFSSKTYQ HDLALLRLRG  
RKNSCAILSP HVQPVCLPSS AAPPSETVLC EVAGWGHQFE GAE EYATFLQ  
EAQVPFISLD RCSSSNVHGD AILPGMLCAG FLEGGADACQ GDSGGPLVCD  
EGVTERQLTL RGVISWGS GC GDRNKPGVYT DVANYLDW IQ EHTAF
```

USAGE

Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.