

Eukaryotic Vascular Endothelial Growth Factor Receptor 1 (VEGFR1) Instruction Manual

SFPB186Hu62

Homo sapiens (Human)

| | |
|---------------------------------|--|
| Source | Eukaryotic expression |
| Host | 293F cell |
| Endotoxin Level | <1.0EU per 1µg (determined by the LAL method) |
| Subcellular Location | Secreted |
| Predicted Molecular Mass | 35.0kDa |
| Accurate Molecular Mass | 50kDa(Analysis of differences refer to the manual) |
| Residues & Tags | Ser27~Ile328 with N-terminal His Tag |
| Buffer Formulation | PBS, pH7.4, containing 5% Trehalose. |
| Traits | Freeze-dried powder |
| Purity | > 95% |
| Isoelectric Point | 9.6 |
| Applications | Positive Control; Immunogen; SDS-PAGE; WB. |

SEQUENCE

```
SKLK DPESLKGQTQ HIMQAGQTLH
LQCRGEEAAHK WSLPEMVSKE SERLSITKSA CGRNGKQFCS TLTLNNTAQAN
HTGFYSCKYL AVPTSKKKET ESAIYIFISD TGRPFVEMYS EIPEIIHMTE
GRELVIPCRV TSPNITVTLK KFPLDTLIPD GKRIIWDSRK GFIISNATYK
EIGLLTCEAT VNGHLYKTNY LTHRQTNTII DVQISTPRPV KLLRGHTLVL
NCTATTPLNT RVQMTWSYPD EKNKRASVRR RIDQSNSHAN IFYSVLTIDK
MQNKDKGLYT CRVRSGPSFK SVNTSVHI
```

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.