Eukaryotic Lysyl Oxidase (LOX) Instruction Manual

SFPC307Hu61

Homo sapiens (Human)

Source	Eukaryotic expression
Host	293F cell
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Secreted
Predicted Molecular Mass	30.6kDa
Accurate Molecular Mass	33kDa(Analysis of differences refer to the manual)
Residues & Tags	Asp169~Tyr417 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	6.3
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

DD PYNPYKYSDD NPYYNYYDTY ERPRPGGRYR PGYGTGYFQY GLPDLVADPY YIQASTYVQK MSMYNLRCAA EENCLASTAY RADVRDYDHR VLLRFPQRVK NQGTSDFLPS RPRYSWEWHS CHQHYHSMDE FSHYDLLDAN TQRRVAEGHK ASFCLEDTSC DYGYHRRFAC TAHTQGLSPG CYDTYGADID CQWIDITDVK PGNYILKVSV NPSYLVPESD YTNNVVRCDI RYTGHHAYAS GCTISPY

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at - 80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

kDa 70 44 33 26 22 18 14 10

Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

Image