Eukaryotic Kynureninase (KYNU) Instruction Manual

SFPD660Hu41

Homo sapiens (Human)

Source Eukaryotic expression

Host 293F cell

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Cytoplasm **Predicted Molecular Mass** 55.5kDa

Accurate Molecular Mass 56kDa(Analysis of differences refer to the manual)

Residues & Tags Met1~Asn465 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 95% Isoelectric Point 7.0

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

MEPSSLELPA DTVQRIAAEL KCHPTDERVA LHLDEEDKLR HFRECFYIPK IQDLPPVDLS LVNKDENAIY FLGNSLGLQP KMVKTYLEEE LDKWAKIAAY GHEVGKRPWI TGDESIVGLM KDIVGANEKE IALMNALTVN LHLLMLSFFK PTPKRYKILL EAKAFPSDHY AIESQLQLHG LNIEESMRMI KPREGEETLR IEDILEVIEK EGDSIAVILF SGVHFYTGQH FNIPAITKAG QAKGCYVGFD LAHAVGNVEL YLHDWGVDFA CWCSYKYLNA GAGGIAGAFI HEKHAHTIKP ALVGWFGHEL STRFKMDNKL QLIPGVCGFR ISNPPILLVC SLHASLEIFK QATMKALRKK SVLLTGYLEY LIKHNYGKDK AATKKPVVNI ITPSHVEERG CQLTITFSVP NKDVFQELEK RGVVCDKRNP NGIRVAPVPL YNSFHDVYKF TNLLTSILDS AETKN

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.