

# Eukaryotic Ephrin B2 (EFNB2) Instruction Manual

## SFPE072Mu61

### Mus musculus (Mouse)

<b>Source</b>	Eukaryotic expression
<b>Host</b>	293F cell
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Secreted
<b>Predicted Molecular Mass</b>	48.8kDa
<b>Accurate Molecular Mass</b>	60kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Arg29~Glu227 with N-terminal His Tag and C-terminal Fc Region of Human IgG1
<b>Buffer Formulation</b>	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 95%
<b>Isoelectric Point</b>	7.9
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

### SEQUENCE

```
RS IVLEPIYWNS SNSKFLPGQG  
LVLYPQIGDK LDIICPKVDS KTVGQYEEYK VYMVDKDQAD RCTIKKENTP  
LLNCARPDQD VKFTIKFQEF SPNLWGLEFQ KNKDYYIIST SNGSLEGLDN  
QEGGVCQTRA MKILMKVGQD ASSAGSARNH GPTRRPELEA GTNGRSSTTS  
PFVKPNPGSS TDGNSAGHSG NLLGSE
```

### USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.  
Do not vortex.

### STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## **Image**




Figure. SDS-PAGE

## **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.